

<p>Drug Facts</p> <p>Active ingredient (in each caplet) Ephedrine sulfate 25 mg.....Bronchodilator</p> <p>Purpose</p> <p>Uses</p> <ul style="list-style-type: none"> ■ for temporary relief of mild symptoms of intermittent asthma: <ul style="list-style-type: none"> ■ wheezing ■ tightness of chest ■ shortness of breath <p>Warnings</p> <p>Asthma alert: Because asthma may be life threatening, see a doctor if you</p> <ul style="list-style-type: none"> ■ are not better in 60 minutes ■ get worse ■ need more than 6 caplets in 24 hours 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ use more than 4 caplets in 24 hours for 3 or more days a week ■ have more than 2 asthma attacks in a week <p>These may be signs that your asthma is getting worse.</p> <ul style="list-style-type: none"> ■ This product will not give you asthma relief as quickly as an inhaled bronchodilator. <p>Do not use</p> <ul style="list-style-type: none"> ■ unless a doctor said you have asthma ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients
<p>Drug Facts (continued)</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ ever been hospitalized for asthma ■ heart disease ■ high blood pressure ■ diabetes ■ thyroid disease ■ seizures ■ narrow angle glaucoma ■ a psychiatric or emotional condition ■ trouble urinating due to an enlarged prostate gland <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ■ taking prescription drugs for asthma, obesity, weight control, depression, or psychiatric or emotional conditions ■ taking any drug that contains phenylephrine, pseudoephedrine, ephedrine, or caffeine (such as for allergy, cough-cold, or pain) <p>When using this product</p> <ul style="list-style-type: none"> ■ your blood pressure or heart rate may go up. This could increase your risk of heart attack or stroke, which may cause death. ■ your risk of heart attack or stroke increases if you: <ul style="list-style-type: none"> ■ have a history of high blood pressure or heart disease ■ take this product more frequently or take more than the recommended dose ■ avoid foods or beverages that contain caffeine 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ avoid dietary supplements containing ingredients reported or claimed to have a stimulant effect <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ your asthma is getting worse (see Asthma Alert) ■ you have difficulty sleeping ■ you have a rapid heart beat ■ you have tremors, nervousness, or seizure <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than directed ■ adults and children 12 years and over: take 1 caplet every 4 hours as needed. Do not take more than 6 caplets in 24 hours or as directed by a doctor. ■ children under 12 years of age: do not use <p>Other information</p> <ul style="list-style-type: none"> ■ store at 20-25°C (68-77°F)
<p>Drug Facts (continued)</p> <p>Inactive ingredients dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, silicon dioxide, sodium starch glycolate, stearic acid, talc</p>	<p>Drug Facts (continued)</p> <p>Questions or comments?</p> <p>call 1-855-565-9800 weekdays 9 AM to 5 PM EST</p>